



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

guided the writers in their selection of material for illustration is not clear. As a list of parasitic fungi from a region where little systematic collection has been carried on, this paper is a useful contribution. Its usefulness might have been greatly enhanced if the authors had indicated what part of their material constituted additions to the known fungus flora of Texas, and what part represented species formerly known from that region, for it can scarcely be doubted that so extensive a collection of material contains much that is new to the region.—H. HASSELBRING.

A new type of Cycadofilicales.—From a study of numerous casts, SCHUSTER²⁵ has described the staminate and ovulate flowers of *Schuetzia anomala* and has drawn some conclusions in regard to the position of the genus. The impressions of the staminate flowers were so numerous that it was not difficult to make reconstructions. The flower consists of 12–20 cyclic sporophylls united throughout the lower two-thirds of their length and bearing sporangia upon their inner surfaces, resembling SELLARD's *Codonotheca* and STUR's *Calymmatotheca*. The flowers are in a spicate inflorescence. The longitudinally striated "seeds" described by GOEPFERT are regarded as megasporophylls, and it is important to note that these megasporophylls are in undoubted connection with twigs bearing conifer-like leaves. On account of this association, SCHUSTER would make *Schuetzia* the type of a new group of Cycadofilicales, characterized by the conifer-like leaves.—CHARLES J. CHAMBERLAIN.

Root nodules of Podocarpaceae.—Miss SPRATT²⁶ has found that root nodules are present in *Podocarpus*, *Microcachrys*, *Dacrydium*, *Saxegothaea*, and *Phyllocladus*, being modified lateral roots. A root-hair is penetrated by *Pseudomonas radicola* (a nitrogen-fixing organism) and from thence enters the cortex. In all cases the nodules are produced by the infection of the meristematic tissue of the young lateral root before it emerges from the cortex of the parent root. Many interesting observations are made upon the stages of the bacteria and also upon the condition of the tissues of the host. The conclusion is suggested that the morphology of the nodules favors the view that *Podocarpus* and *Saxegothaea* "are the most widely divergent of the genera in the Podocarpaceae, and that they are connected through *Microcachrys* and *Dacrydium*." The presence of the nodules in *Phyllocladus* is also further confirmation that the genus is related to the podocarps rather than to the taxads.—J. M. C.

²⁵ SCHUSTER, J., Über die Fruktifikation von *Schuetzia anomala*. Sitzungsber. Kaiserl. Akad. Wiss. Wien 120:1125–1134. pls. 1, 2. 1911.

²⁶ SPRATT, ETHEL ROSE, The formation and physiological significance of root nodules in the Podocarpaceae. Ann. Botany 26:801–814. pls. 77–80. 1912.